

Characterization of proteases isolated from *Kudoa septempunctata*

Sang Phil Shin^{1,*}, Kosuke Zenke², Hiroshi Yokoyama²

¹Fisheries Laboratory, Kinki University, Wakayama 649-2211, Japan

²Department of Aquatic Bioscience, Graduate School of Agricultural and Life Sciences, University of Tokyo, Tokyo 113-8657, Japan

(Received: July 28, 2015; Revised: August 9, 2015; Accepted: August 17, 2015)

Abstract : Proteases play important roles in parasite development and host parasite interactions. The protease of *Kudoa* spp. has been recognized as a key factor of severe proteolysis of fish muscle post-mortem; however, there is little information available regarding the protease of *Kudoa* (*K.*) *septempunctata*, which was recently identified as a cause of food poisoning in humans. The present study was conducted to isolate and characterize proteases to elucidate the type of protease contained in the parasite and determine the optimal pH for protease activity. We confirmed the cysteine protease and metalloprotease produced by *K. septempunctata*. While the cysteine protease showed optimal activity at pH 5 that decreased rapidly with increasing pH, the optimal activity of metalloprotease was pH 7, and it remained stable from pH 6 to pH 8. These results indicate that the pH of cysteine protease is not proper for fish muscle post-mortem, and that metalloprotease can act in human intestines. Overall, the present study provides important information that improves our understanding of the role of protease physiology and the subsequent food poisoning caused by *K. septempunctata*.

Keywords : pH, cysteine protease, *Kudoa septempunctata*, metalloprotease

Introduction

The family *Kudoidae* (Myxozoa: *Multivalvulida*) consist of a single genus *Kudoa* Meglitsch, 1947, having four or more shell valves and polar capsules, and more than ninety species of *Kudoa* have been described from marine and estuarine fish species [10, 27]. Most of *Kudoa* species are histozoic, inducing macroscopic cysts in various organs including somatic muscles, however some species forming a pseudocyst in the myofibers and cause post-mortem myoliquefaction [18, 22]. Although myxozoan parasites are generally considered harmless to humans, certain human illnesses have been attributed to *Kudoa* sp. such as allergic symptoms [14] and food-borne diarrhea. Serial food poisonings associated with ingestion of raw olive flounder *Paralichthys olivaceus* have been recently reported in Japan, and epidemiological analysis demonstrates that *Kudoa* (*K.*) *septempunctata* is associated with these illnesses [11].

Proteases play an integral role in interactions between parasites and their hosts, are involved in parasite physiology development [3, 19-21], and contribute to parasite virulence [12, 24]. These enzymes have been also described in some of *Kudoa* species such as *K. rosenbuschi*, *K. paniformis*, and *K. thyrsites*. They are associated with a degradation of host muscle tissue post-mortem, which was known as myolique-

faction [7, 15]. However, there is currently no information about the proteases of *K. septempunctata* even the parasite has threatened a public health.

The objective of this study was to isolate and characterize the parasite proteases that may contribute to the physiology and pathogenicity of *K. septempunctata*. We focused on whether *K. septempunctata* possesses what kind of proteases, and what of pH does work on an optimal for the activity of proteases. The information in the present study will contribute to understand the life cycle or a relationship between the role of protease and food poisoning of *K. septempunctata*.

Materials and Methods

Sample preparation

Olive flounder samples suspected of being infected with *K. septempunctata* were obtained from an olive flounder farm in Japan. The fish samples were examined microscopically to confirm the presence of *K. septempunctata* spores in squash preparations. The spores were purified using a modified Percoll (Sigma-Aldrich, USA) density-gradient centrifugation method as previously described [26]. Purified spores were diluted in phosphate buffer solution (PBS) to a final concentration of 10^7 spores/ml. The lysate of *K. septempunctata*

*Corresponding author

Tel: +81-739-42-2625, Fax: +81-739-42-2634

E-mail: charisma-shin@daum.net

(LKs) was obtained as described by previous report [25], with some modifications. Briefly, the spores were washed twice by centrifugation at $10,000 \times g$ for 2 min in 0.25 M sucrose. The spores were then disrupted ultrasonically. The homogenate was centrifuged at $15,000 \times g$ for 10 min and the resulting supernatant fraction was stored in aliquots at -80°C . Protein concentration in the extract was determined by using the Pierce BCA protein assay kit (Thermo Fisher Scientific, USA) with a microplate reader (MPRA4; Tosoh, Japan) at 570 nm. The bovine serum albumin was used as a standard.

Affinity chromatography

The LKs was dialyzed at 4°C against four volumes of 25 mM acetate buffer (pH 5.0) and bacitracin was used in affinity chromatography as recommended [4] with some modifications. Bacitracin was coupled to cyanogen bromide-activated Sepharose 4B (Sigma-Aldrich) and the dialyzed lysate was applied to a 5×1 cm column of bacitracin-Sepharose pre-equilibrated with buffer A (25 mM acetate buffer, pH 5.0), at a flow rate of 0.5 ml/min. The column was washed with 50 ml of buffer A (0.5 mL/min) and eluted with buffer B (buffer A containing 1 M NaCl, 25% of isopropanol) at the same flow rate. The eluted fractions were pooled in conical tubes and concentrated using ultrafiltration through Amicon Ultra-15 filter (30-kDa cutoff; Millipore, USA), and desalted twice using Tris-HCl buffer (TBS; pH 7.0). The partial purified protease of *K. septempunctata* (PKs) was concentrated again using Amicon Ultra-0.5mL centrifuga filter and the concentration of protein was determined as described above.

SDS-PAGE and gelatin-zymography

The LKs was analyzed by sodium dodecyl sulfate-polyacrylamide gel electrophoresis (SDS-PAGE). Sample (20 μg) was mixed with SDS-PAGE loading buffer and electrophoresis was carried out for 100 min in a Tris-glycine buffer system. The gel was stained with Coomassie blue for the visualization of the proteins, and molecular weights estimated by comparison with SDS-PAGE protein standards. In order to visualize the proteases activity of LKs, gelatin was added to the 10% acrylamide running gel for a final concentration of 0.2% (w/v) protein. The electrophoresis of two lysate

samples (each 3 μg) was carried out as condition described above. After electrophoresis the gels were incubated for 30 min in 2.5% (v/v) Triton X-100 at room temperature, and then each of gels were incubated for 18 h in 50 mM 2-N-morpholino ethanesulfonic acid (MES) buffer (200 mM NaCl, 0.02% Brij-35 and 0.01% NaN_3 , pH 5.0) and TBS (200 mM NaCl, 0.02% Brij-35 and 0.01% NaN_3 , pH 7.0) at 37°C , respectively. The gels were stained with Coomassie blue, and destained until clear gel bands were apparent where proteolytic hydrolysis of the gel embedded substrate had occurred. In addition, we also examined the activity of PKs through gelatin-zymography described above and it was developed in TBS (200 mM NaCl, 0.02% Brij-35 and 0.01% NaN_3 , pH 7.0).

Protease activity and inhibitor assays using azocasein

An azocasein assay was used to further characterize the activity of the LKs and PKs. Protease activity was determined by measuring the release of acid-soluble material from azocasein (Sigma-Aldrich). The optimum pH for both the LKs and PKs was determined within a pH range of 4.0 to 8.0, using 50 mM sodium acetate (pH 4.0–6.0) and 50 mM Tris-HCl (pH 7.0–8.0) buffers. The reaction was initiated by the addition of 10- μL samples to 90 of μL reaction buffer containing 50 mM sodium acetate/Tris-HCl buffer, pH 4.0–8.0, 0.2% azocasein, 200 mM NaCl, 0.02% Brij-35, 0.01% NaN_3 , and the mixtures were incubated for 18 h at 37°C . The reaction was stopped by the addition of 100 μL 10% (w/v) trichloroacetic acid (TCA), and the mixtures were incubated at room temperature for 30 min. The mixtures were then centrifuged ($12,000 \times g$, 10 min) to pellet the unreacted substrate, and 125 μL of the supernatant (containing the TCA-soluble azo-compounds) was withdrawn and mixed with 50 μL 2.0 M NaOH. The absorbance of the mixture was read at 440 nm by MPRA4. The effects of various inhibitors (Pepstatin A, PMSF, EDTA, 1, 10-phenanthroline, E64, and leupeptin) against the LKs and PKs were evaluated in sodium acetate buffer (pH 5.0) and TBS (pH 7.0), respectively. The final concentrations of inhibitors indicated in Table 1 and the results reported from all assays were carried out in triplicate.

Table 1. Inhibition assay of proteases by various inhibitors

Inhibitor	Specificity	Concentration (mM)	Lysate proteases activity (% control)*	Partial purified protease activity (% control)*
E64	Cystein proteases	0.005	3 ± 1	98 ± 4
Leupeptin	Cystein proteases and trypsin like serine proteases	0.05	4 ± 2	96 ± 5
PMSF	Serine protease	0.5	99 ± 3	97 ± 4
Pepstatin	Asparatic proteases	0.005	99 ± 1	98 ± 4
1,10-Phenanthroline	Metalloproteases	5	82 ± 2	1 ± 1
EDTA	Metalloproteases	5	108 ± 2	2 ± 1

*Enzyme activity was estimated using azocasein as a substrate. The results are mean values from triplicate experiments \pm SD.




Fig. 1. Analysis of the lysate proteases (LKs) and partial purified protease (PKs) of *Kudoa (K.) septempunctata* by 10% sodium dodecyl sulfate-polyacrylamide and gelatin zymography. Bands were visualized by Coomassie Blue staining (lane 1) and destaining (lanes 2–4). Lane M, molecular mass protein markers; lane 1, crude supernatant from homogenized purified *K. septempunctata*; lane 2, LKs was incubated at MES buffer (pH 5.0); lane 3, LKs was incubated at Tris-HCl buffer (TBS; pH 7.0); lane 4, PKs was incubated at TBS (pH 7.0).




Fig. 2. pH activity profile of LKs and PKs. The activity of the proteases against azocasein substrate was measured at different pHs. The activities are shown relative to that at pH 5.0 and pH 7.0 being 100%, respectively. The results are mean values from triplicate experiments \pm SD.

Results

The LKs showed bands of various sizes by SDS-PAGE. Six of the bands (83, 75, 63, 48, 29, and 26.5 kDa) exhibited gelatinolytic activity at pH 5.0 (lanes 1 and 2 in Fig. 1). Only one of the six LKs bands (83 kDa) had detectable protease activity at pH 7.0 (lane 3 in Fig. 1), which was larger than the only PKs band (45 kDa) that exhibited gelatinolytic activity at pH 7.0 (lane 4 in Fig. 1). The LKs was enzymatically active against the azocasein at pH 5.0, with activity that reduced dramatically at pH 6.0 (Fig. 2A). The PKs showed optimal activity at pH 7.0 and was stable between pH 6.0–8.0 (Fig. 2B). While proteolytic activity of LKs was largely inhibited by adding the cysteine protease specific inhibitors E64 and leupeptin, the PKs was inhibited by metalloprotease

inhibitors such as 1,10-phenanthroline and EDTA. The effect of the inhibitors on protease activity of LKs and PKs using azocasein as the substrate are shown in Table 1.

Discussion

Proteases play important roles in interactions between parasites and their hosts, and underlie the pathogenicity of many organisms [19, 21]. The cysteine proteases have been described in *Kudoa* spp. and are possibly a key factor of the parasites, which causes post-mortem myoliquefaction [7, 15]. Although *K. septempunctata* is known to cause food poisoning in humans when accidentally consumed with raw flounder, there are poorly understood about the proteases of this parasite and the *K. septempunctata* infected fish muscle has showed no myoliquefaction [11, 18]. The present study identified a presence of cysteine proteases through LKs that showed the optimal enzymatic activity at pH 5.0. However, the activity of cysteine protease was decreased rapidly between pH 5.0 and pH 6.0. Following death of the fish, the stored carbohydrate glycogen is anaerobically degraded and lactic acid is accumulated in muscle resulting in a pH drop from 7.4 to 6.0 [2]. In addition, previous studies have reported that the pH of muscle in post-mortem flounder and halibut decreases to pH 6.5 for 3 days [6, 16]. Therefore, we suggest that *K. septempunctata* does not cause post-mortem myoliquefaction based on the relationship between the optimal pH of the parasite cysteine protease and the pH of muscle in post-mortem fish.

In previous study, we reported that factors affecting sporoplasm release in *K. septempunctata* such as enzymes, culture media, and protease inhibitors. Interestingly, the sporoplasm release was prevented by 1,10-phenanthroline (metalloprotease) and it suggested that metalloprotease is related the sporoplasm release in *K. septempunctata* [26]. Metalloproteases of parasite have been related to pathogenesis and are involved in processes such as immunity evasion, development, and metabolism [8, 17]. In addition, the activity of metalloproteases are affected by pH of environment [13]. In the present study, we observed one protease (83 kDa) of LKs which showed the proteolytic activity at pH 7.0 and that was purified by affinity chromatography partially. We speculate that the protease was processed as a mature form (45 kDa) for purifying. The PKs have optimal activity at pH 7.0 and showed stable activity from pH 6.0 to pH 8.0. However, the activity was completely prevented by inhibitors of metalloprotease. These results demonstrated that the PKs is a metalloprotease which has activity from pH 6.0 to pH 8.0. We also observed the prevention of sporoplasm release in acidified culture medium [26], even there are the factors to release it. It is indicated that the environmental pH for *K. septempunctata* is a key factor to release sporoplasm. Human intraluminal pH is rapidly changed from highly acidic in the stomach to about pH 6.0 in the duodenum, and the pH gradually increases in the small intestine from pH 6.0 to about pH 7.4

in the terminal ileum. The pH drops to 5.7 in the caecum, but again gradually increases, reaching pH 6.7 in the rectum [5]. Therefore, the pH of human intestine is proper to work for the PKs especially that is optimal pH in small intestine. The median latency between eating the *K. septempunctata* infected raw fish and the presence of the first symptoms is five hours that is corresponding with the time of dietary transit reach to small intestine [1, 9, 28]. Based on these results, we suggest that the PKs contribute the sporoplasm in small intestine, which causes food poisoning by *K. septempunctata*. Two of possible ways to contribute the sporoplasm release were speculated: 1) PKs effect on the mechanism of actin polymerization which discharges the sporoplasm by condensing actin filament [23], 2) the metalloprotease is involved in metabolism of glucose related to the release of sporoplasm in *K. septempunctata* [26]. In addition, we also observed the proteolytic activity (5–7%) in LKs at pH 6–8, suggesting that the activity was produced by the metalloprotease and the protease is not related to post-mortem myoliquefaction.

In conclusion, the present study demonstrates the presence and kind of proteases in *K. septempunctata*. The cysteine protease showed the strong activity at pH 5 although it is decreased rapidly at pH 6 over while metalloprotease have stable activity from pH 6 to 8. These findings elucidate the relation between proper pH range of the protease and pathophysiological response such as post-mortem myoliquefaction and sporoplasm release in *K. septempunctata*. The information of present study will contribute to reveal the life cycle of *K. septempunctata* and establish strategies to treat and prevent disease caused by this parasite. However, further studies will be required to demonstrate the molecular and immunological mechanism of the proteases effect on physiological response in life cycle of *K. septempunctata*.

Acknowledgments

This research was supported by International Research Fellow of the Japan Society for the Promotion of Science (P14398; grant no. 26-04398) and Basic Science Research Program through the National Research Foundation of Korea (NRF) funded by the Ministry of Education, Science and Technology (2013R1A6A3A03063215).

References

1. **Degen LP, Phillips SF.** Variability of gastrointestinal transit in healthy women and men. *Gut* 1996, **39**, 299-305.
2. **Delbarre-Ladrat C, Chéret R, Taylor R, Verrez-Bagnis V.** Trends in postmortem aging in fish: understanding of proteolysis and disorganization of the myofibrillar structure. *Crit Rev Food Sci Nutr* 2006, **46**, 409-421.
3. **Dörfler C, El-Matbouli M.** Isolation of a subtilisin-like serine protease gene (*MlxSubtSP*) from spores of *Myxobolus cerebralis*, the causative agent of whirling disease. *Dis Aquat Organ* 2007, **73**, 245-251.
4. **Faisal M, Schafhauser DY, Garreis KA, Elsayed E, La Peyre JF.** Isolation and characterization of *Perkinsus marinus* proteases using bacitracin-sepharose affinity chromatography. *Comp Biochem Physiol B Biochem Mol Biol* 1999, **123**, 417-426.
5. **Fallingborg J.** Intraluminal pH of the human gastrointestinal tract. *Dan Med Bull* 1999, **46**, 183-196.
6. **Foy RJ, Crapo CA, Kramer DE.** Investigating the roles of temperature and exercise in the development of chalkiness in Pacific halibut. Report No. 50. International Pacific Halibut Commission (US), Washington, 2006.
7. **Funk VA, Olafson RW, Raap M, Smith D, Aitken L, Haddow JD, Wang D, Dawson-Coates JA, Burke RD, Miller KM.** Identification, characterization and deduced amino acid sequence of the dominant protease from *Kudoa paniformis* and *K. thyrssites*: a unique cytoplasmic cysteine protease. *Comp Biochem Physiol B Biochem Mol Biol* 2008, **149**, 477-489.
8. **Gras S, Byzia A, Gilbert FB, McGowan S, Drag M, Silvestre A, Niepceon A, Lecaille F, Lalmanach G, Brossier F.** Aminopeptidase N1 (EtAPN1), an M1 metalloprotease of the apicomplexan parasite *Eimeria tenella*, participates in parasite development. *Eukaryot Cell* 2014, **13**, 884-895.
9. **Iwashita Y, Kamijo Y, Nakahashi S, Shindo A, Yokoyama K, Yamamoto A, Omori Y, Ishikura K, Fujioka M, Hatada T, Takeda T, Maruyama K, Imai H.** Food poisoning associated with *Kudoa septempunctata*. *J Emerg Med* 2013, **44**, 943-945.
10. **Jeon CH, Wi S, Song JY, Choi HS, Kim JH.** Development of loop-mediated isothermal amplification method for detection of *Kudoa septempunctata* (Myxozoa: Multivalvulida) in olive flounder (*Paralichthys olivaceus*). *Parasitol Res* 2014, **113**, 1759-1767.
11. **Kawai T, Sekizuka T, Yahata Y, Kuroda M, Kumeda Y, Iijima Y, Kamata Y, Sugita-Konishi Y, Ohnishi T.** Identification of *Kudoa septempunctata* as the causative agent of novel food poisoning outbreaks in Japan by consumption of *Paralichthys olivaceus* in raw fish. *Clin Infect Dis* 2012, **54**, 1046-1052.
12. **Mackey ZB, O'Brien TC, Greenbaum DC, Blank RB, McKerrow JH.** A cathepsin B-like protease is required for host protein degradation in *Trypanosoma brucei*. *J Biol Chem* 2004, **279**, 48426-48433.
13. **Malagón D, Adroher FJ, Díaz-López M, Benítez R.** Collagenolytic activity related to metalloproteases (and serine proteases) in the fish parasite *Hysterothylacium aduncum* (Nematoda: Anisakidae). *Dis Aquat Organ* 2010, **90**, 129-134.
14. **Martínez de Velasco G, Rodero M, Cuéllar C, Chivato T, Mateos JM, Laguna R.** Skin prick test of *Kudoa* sp. antigens in patients with gastrointestinal and/or allergic symptoms related to fish ingestion. *Parasitol Res* 2008, **103**, 713-715.
15. **Martone CB, Spivak E, Busconi L, Folco EJE, Sánchez JJ.** A cysteine protease from myxosporean degrades host myofibrils in vitro. *Comp Biochem Physiol B Biochem Mol Biol* 1999, **123**, 267-272.
16. **Massa AE, Palacios DL, Paredi ME, Crupkin M.** Postmortem changes in quality indices of ice-stored flounder (*Paralichthys patagonicus*). *J Food Biochem* 2005, **29**, 570-590.

17. **Matheoud D, Moradin N, Bellemare-Pelletier A, Shio MT, Hong WJ, Olivier M, Gagnon E, Desjardins M, Descoteaux A.** *Leishmania* evades host immunity by inhibiting antigen cross-presentation through direct cleavage of the SNARE VAMP8. *Cell Host Microbe* 2013, **14**, 15-25.
18. **Matsukane Y, Sato H, Tanaka S, Kamata Y, Sugita-Konishi Y.** *Kudoa septempunctata* n. sp. (Myxosporea: Multivalvulida) from an aquacultured olive flounder (*Paralichthys olivaceus*) imported from Korea. *Parasitol Res* 2010, **107**, 865-872.
19. **McKerrow JH.** Parasite proteases. *Exp Parasitol* 1989, **68**, 111-115.
20. **McKerrow JH.** Development of cysteine protease inhibitors as chemotherapy for parasitic diseases: insights on safety, target validation, and mechanism of action. *Int J Parasitol* 1999, **29**, 833-837.
21. **McKerrow JH, Sun E, Rosenthal PJ, Bouvier J.** The proteases and pathogenicity of parasitic protozoa. *Annu Rev Microbiol* 1993, **47**, 821-853.
22. **Moran JDW, Whitaker DJ, Kent ML.** A review of the myxosporean genus *Kudoa* Meglitsch, 1947, and its impact on the international aquaculture industry and commercial fisheries. *Aquaculture* 1999, **172**, 163-196.
23. **Ohnishi T, Kikuchi Y, Furusawa H, Kamata Y, Sugita-Konishi Y.** *Kudoa septempunctata* invasion increases the permeability of human intestinal epithelial monolayer. *Foodborne Pathog Dis* 2013, **10**, 137-142.
24. **Que X, Reed SL.** Cysteine proteinases and the pathogenesis of amebiasis. *Clin Microbiol Rev* 2000, **13**, 196-206.
25. **Shin SP, Han SY, Han JE, Jun JW, Kim JH, Park SC.** Expression and characterization of cathepsin L-like cysteine protease from *Philasterides dicentrarchi*. *Parasitol Int* 2014, **63**, 359-365.
26. **Shin SP, Zenke K, Yokoyama H, Yoshinaga T.** Factors affecting sporoplasm release in *Kudoa septempunctata*. *Parasitol Res* 2015, **114**, 795-799.
27. **Whipps CM, Gossel G, Adlard RD, Yokoyama H, Bryant MS, Munday BL, Kent ML.** Phylogeny of the multivalvulidae (Myxozoa: Myxosporea) based on comparative ribosomal DNA sequence analysis. *J Parasitol* 2004, **90**, 618-622.
28. **Worsøe J, Fynne L, Gregersen T, Schlageter V, Christensen LA, Dahlerup JF, Rijkhoff NJ, Laurberg S, Krogh K.** Gastric transit and small intestinal transit time and motility assessed by a magnet tracking system. *BMC Gastroenterol* 2011, **11**, 145.