

***Corresponding author:**

Seong-Mok Jeong
Department of Veterinary Surgery, College of Veterinary Medicine, Chungnam National University, 99 Daehak-ro, Yuseong-gu, Daejeon 34134, Korea
Tel: +82-42-821-6763
Fax: +82-42-821-6703
E-mail: jsmok@cnu.ac.kr

ORCID:

<https://orcid.org/0000-0001-8653-5927>


Conflict of interest:

The authors declare no conflict of interest.

Received: Nov 10, 2021

Revised: Dec 30, 2021

Accepted: Jan 4, 2022



© 2022 The Korean Society of Veterinary Science.

© This is an open-access article distributed under the terms of the Creative Commons Attribution Non-Commercial license (<http://creativecommons.org/licenses/by-nc/4.0/>), which permits unrestricted non-commercial use, distribution, and reproduction in any medium, provided the original work is properly cited.

Assessment of prognostic factors in dogs with mammary gland tumors: 60 cases (2014-2020)

Chang-Hwan Moon¹, Dae-Hyun Kim¹, Sung-Ho Yun², Hae-Beom Lee¹, Seong-Mok Jeong^{1,7}

¹Department of Veterinary Surgery, College of Veterinary Medicine, Chungnam National University, Daejeon 34134, Korea

²Department of Veterinary Surgery, College of Veterinary Medicine, Kyungpook National University, Daegu 41566, Korea

Abstract

Canine mammary gland tumors are the most common neoplasms in intact female dogs. Approximately half of all mammary tumors are malignant, and there is a risk of metastasis, which is associated with a poor prognosis. This study was to evaluate the prognostic factors of canine mammary gland tumors and the risk factors associated with the development of malignant tumors. From 2014 to 2020, 60 dogs with mammary gland tumors that underwent surgical treatment were evaluated in this retrospective study. Tumor size, TNM stage, and histopathological results were prognostic factors for 2-year survival after surgery. Every 10 mm increase in tumor size, increased the risk of death within 2 years after surgery 1.213 times. Dogs with TNM stage IV or V had 8.667 fold risk of death within 2 years after surgery. The 2-year survival rate for dogs with benign tumors was 90.2% and for malignant tumors was 67.3%. Tumor size is the most important prognostic factor for canine mammary gland tumors. As tumor size increased by 10 mm, the risk for development of malignant tumors increased by 1.487 times. Tumors larger than 30 mm are highly likely to be malignant, and metastatic evaluation and wide resection should be considered.

Keywords: dogs; mammary gland tumor; risk factors; tumor size

Introduction

Mammary gland tumors (MGTs) are the most common neoplasm in intact female dogs that account for over 40% of all tumors [1,2]. Surgical resection remains the standard treatment for most types of MGTs, except in cases of inoperable highly metastatic disease and most inflammatory mammary carcinomas [1,3,4]. Surgical efficacy and feasibility depend on the number of tumors, tumor size, location, and patient status [5]. Dogs in early stages of tumor progression, according to the TNM classification of malignant tumors, and small tumors are often treated effectively with surgery alone. Dogs with benign MGTs and approximately 50% of dogs with malignant MGTs are also treated with surgery alone [6,7].

The risk factors that influence the prognosis of dogs with MGTs include age, ovariectomy (OHE) at the time of tumor resection, tumor size, TNM stage, and tumor histopathological type. Increased age has been associated with lower 2-year survival rates in dogs with MGTs [8]. It has been reported that there is no

prolonged survival time in dogs with OHE at the time of tumor resection, compared with dogs that were not spayed [9-11]; however, dogs treated with OHE at the time of tumor resection lived significantly longer than dogs treated with tumor removal alone [12]. Tumor size is one of the most important risk factors associated with the prognosis of dogs with MGTs [13-16]. Dogs with a low-level TNM stage lived significantly longer than dogs in late stage [15-18]. Metastasis to regional lymph nodes, distant metastasis, and TNM stage are strongly associated. Approximately 35% to 50% of canine MGTs are malignant [4] and most malignant tumors eventually metastasize [19]. The likelihood of metastasis increases when the tumor is malignant [17,20].

Preoperative evaluation of prognosis provides the veterinarian with important information that may influence treatment planning. The purpose of this study was to evaluate the prognostic factors of MGTs in dogs and to suggest appropriate guidelines for treatment.

Materials and Methods

Study population and inclusion criteria

The medical records of dogs with MGTs that received surgical treatment at the Chungnam National University Veterinary Medical Teaching Hospital from 2014 to 2020 were reviewed and included in this study. Dogs without preoperative examination results, histological examination results, or follow-up data were excluded from this study. Data obtained from the medical records, including clinical data such as signalment, blood analysis results (complete blood count, serum biochemical profile), and cytological examination results (fine needle aspiration), were included. Tumor number, size, and location were also included. Tumor size was decided as the maximum diameter of the tumor among all of the MGTs for each dog. Thoracic radiography and standard abdominal ultrasonography were performed. Computed tomography was performed in patients with malignant tumors. Tumor resection was performed based on the diagnostic examination, and surgical techniques (lumpectomy, simple mastectomy, regional mastectomy, unilateral mastectomy or total mastectomy) were recorded. Histopathological examination was performed after the tumor resection. The clinical stage of the MGTs was assigned according to a modified version of the World Health Organization TNM staging classification [3,21].

Follow-up examination

Follow-up examinations, including physical examination and imaging procedures (thoracic radiography and abdominal so-

nography), were performed every 3 months after surgery. When follow-up examinations were not possible, data were obtained through telephone interviews with the owners. Follow-up data included local relapse, distant metastasis, and MGT-related death. The dogs died by metastasis, recurrence or euthanatized for MGT was defined as MGT-related death. In dogs that died of causes unrelated to the MGTs, death was defined as censored data for calculating survival time.

Statistical analysis

The data obtained included sex, age, body weight, number of tumors, tumor size, TNM stage, histopathological results, surgical procedures, ovario-uterine disease, relapse, and death. Overall survival (OS) and disease-free period (DFP) were evaluated. OS was calculated from the date of surgical removal of the tumor to the date of death or the last follow-up date. The DFP was defined as the period from tumor resection to relapse. Kaplan-Meier survival curve analysis was used to create category-specific survival curves. The Cox proportional hazards model was used to estimate the effect of each factor (sex, age, body weight, the number of tumors, tumor size, lymph node involvement, distant metastasis, TNM stage, histopathological results, surgical procedures, ovario-uterine disease, and OHE with tumor resection) potentially associated with survival time after tumor resection. The Mann-Whitney test was used to investigate the differences between benign and malignant tumors depending on their size. Logistic regression was used to estimate the risk factors (sex, age, body weight, number of tumors, tumor size, and ovario-uterine disease) associated with malignant MGTs. All statistical analyzes were performed using commercial statistical software (IBM SPSS Statistics ver. 24.0; IBM Corp., USA).

Results

In this study, 69 dogs underwent tumor resection for the surgical treatment of MGTs. Of those, 7 dogs that had no histological examination and 2 dogs that had no follow-up data were excluded. A total of 60 dogs that received surgical treatment for MGTs between 2014 and 2020 met the criteria for inclusion in the study.

The breed distribution was as follows: Beagle (n = 1), Chihuahua (n = 1), Cocker Spaniel (n = 4), Jindo (n = 2), Maltese (n = 15), miniature Pincher (n = 1), miniature Schnauzer (n = 2), mixed (n = 8), papillon (n = 1), Pomeranian (n = 2), Poodle (n = 9), Shih Tzu (n = 7), Spitz (n = 1), Welsh Corgi (n = 1), and Yorkshire Terrier (n = 5).

Of a total of 60 dogs, 51 dogs (85.0%, 51/60) were intact females and 9 dogs (15.0%, 9/60) were spayed females. There were 36 dogs (60.0%, 36/60) with benign tumors (complex adenoma [n = 16], adenoma [n = 10], and benign mixed tumor [n = 10]) and 24 dogs (40.0%, 24/60) with malignant tumors (adenocarcinoma [n = 13], carcinoma [n = 9], extraskeletal osteosarcoma [n = 1], and soft tissue sarcoma [n = 1]). Of the intact female dogs, 32 had benign tumors and 19 had malignant tumors. Of the spayed female dogs, 4 dogs had benign tumors and 5 dogs had malignant tumors.

The incidences of benign and malignant tumors are summarized in Table 1. The mean age of the subjects with benign tumors was 10.9 ± 3.1 years (median, 11.0; range, 3.0-18.0 years) and mean body weight was 5.4 ± 4.2 kg (median, 3.8; range, 2.1-24.0 kg). The mean age for malignant tumors was 11.7 ± 2.9 years (median, 12.0; range, 2.0-16.0 years) and mean body weight was 5.1 ± 2.6 kg (median, 4.6; range, 2.0-11.0 kg).

In this study, there was a significant difference between tumor size and histopathological tumor type ($p = 0.008$). The mean

size \pm standard deviation of benign tumors was 20.0 ± 15.2 mm (median, 15.5 mm; range, 3-60 mm) and for malignant tumors, 48.7 ± 40.2 mm (median, 37.5 mm; range, 7-141 mm). Tumor size was correlated with malignancy ($p = 0.002$).

As tumor size increased by 10 mm, the risk of developing malignant tumors increased 1.487 times ($p = 0.004$). Tumors larger than 50 mm had an 11.815-fold risk of developing into malignant tumors compared to tumors less than 30 mm ($p = 0.001$).

Tumor size, malignancy and high-level TNM stage were significantly associated with 2-year survival after tumor resection (Table 2). Every 10 mm increase in tumor size, increased the risk of death within 2 years after tumor resection 1.213 times ($p = 0.011$). Dogs with tumors larger than 50 mm had a 5.448 higher risk of death within 2 years after tumor resection compared to those with tumors less than 30 mm ($p = 0.021$). In dog with the presence of lymph node involvement, the risk of death within 2 years after tumor resection increased by 8.667 times ($p = 0.001$) compared to those with no lymph node involvement, whereas in dogs with the presence of distant metastasis, the risk was increased 72.665 times ($p = 0.003$) compared to those with no distant metastasis. Dogs with high-level TNM stage (IV or V) were 8.667 times at risk of death within 2 years after tumor

Table 1. Incidence of benign and malignant mammary gland tumors in 60 dogs

Variable	Number of dogs (%)		
	Benign	Malignant	Total
Sex			
Female	32 (62.7)	19 (37.3)	51 (85.0)
Spayed female	4 (44.4)	5 (55.6)	9 (15.0)
Age (y)	10.9 ± 3.1	11.7 ± 2.9	11.2 ± 3.0
Body weight (kg)	5.4 ± 4.2	5.1 ± 2.6	5.3 ± 3.6
Number of tumors			
Solitary	7 (77.8)	2 (22.2)	9 (100.0)
Double	6 (66.7)	3 (33.3)	9 (100.0)
Multiple (≥ 3)	23 (54.8)	19 (45.2)	42 (100.0)
Tumor size (mm)			
T ₁ : < 30	29 (76.3)	9 (23.7)	38 (100.0)
T ₂ : ≥ 30 , < 50	4 (50.0)	4 (50.0)	8 (100.0)
T ₃ : ≥ 50	3 (21.4)	11 (78.6)	14 (100.0)
TNM Stage			
I	28 (82.4)	6 (17.6)	34 (100.0)
II	5 (62.5)	3 (37.5)	8 (100.0)
III	3 (25.0)	9 (75.0)	12 (100.0)
IV	0 (0.0)	5 (100.0)	5 (100.0)
V	0 (0.0)	1 (100.0)	1 (100.0)
Ovario-uterine disease			
Spayed female	4 (44.4)	5 (55.6)	9 (100.0)
Female: present	26 (61.9)	16 (38.1)	42 (100.0)
Female: absent	6 (66.7)	3 (33.3)	9 (100.0)

Values are presented as number (%) or mean \pm standard deviation.

Table 2. Prognostic factors associated with 2-year survival after surgery in 60 dogs with mammary gland tumors

	p-value	Hazard ratio	95% CI
Sex	0.436	0.535	0.111-2.581
Age (y)	0.183	1.176	0.926-1.493
Body weight (kg)	0.704	1.031	0.870-1.221
Number of affected mammary glands	0.212	1.149	0.924-1.428
Number of tumors			
Solitary	0.866	Referent	NA
Double	0.989	50,748.756	0.000-2.007
Multiple (≥ 3)	0.957	89,401.784	0.000-3.521
Tumor size (10 mm)	0.011	1.213	1.046-1.406
Tumor size (T)			
T ₁ : < 30 mm	0.063	Referent	NA
T ₂ : ≥ 30 mm, < 50 mm	0.669	1.644	0.169-15.990
T ₃ : ≥ 50 mm	0.021	5.448	1.295-22.923
Regional lymph node involvement (N)	0.001	8.667	2.319-32.394
Distant metastasis (M)	0.003	72.665	4.453-1,185.538
TNM stage I, II, III vs. IV, V	0.001	8.667	2.319-32.394
Histopathological result	0.022	6.29	1.304-30.351
Ovario-uterine disease	0.748	1.416	0.170-11.790
OHE at tumor resection	0.535	0.535	0.000-986.226

CI, confidence interval; NA, not available; OHE, ovariectomy.

resection compared to those with low-level TNM stage (I, II, or III).

There was a significant difference in the 2-year survival rate between dogs with low-level TNM stage and high-level TNM stage ($p < 0.001$). The 2-year survival rate for dogs with low-level TNM stage was 87.8%, and 33.3% for dogs with high-level TNM stage (Fig. 1A). Mean survival time of dogs with low-level TNM stage was 22.2 ± 0.8 months, but for those with a progressed TNM stage was 11.2 ± 3.6 months.

The 2-year survival rate for dogs with benign tumors was 90.2% and for malignant tumors was 67.3% (Fig. 1B). Mean survival of dogs with benign tumors was 23.1 ± 0.6 months, but 17.7 ± 2.0 months for dogs with malignant tumors. There was a significant difference in the 2-year survival rate after tumor resection between benign and malignant tumors ($p = 0.008$).

Of the total of 60 dogs, 17 dogs (28.3%, 17/60) died within 2 years after tumor resection. Of the 17 dogs, 8 had benign tumors. One dog with a benign tumor died due to the occurrence of inflammatory carcinoma. Another dog died during chemotherapy because of tumor relapse and lymph node metastasis. Six dogs with benign tumors died from unrelated causes of MGTs. Nine dogs with malignant tumors died within 2 years, of which 7 died by MGTs and 2 died from unrelated causes.

Discussion

This study analyzed 60 cases of canine MGTs to evaluate the prognosis after tumor resection. Previous studies revealed that the prognostic factors of dogs with MGTs are age, tumor size, TNM stage, and OHE at the time of tumor resection [6,8-11,13-15,17-18, 22].

Approximately 35% to 50% of canine MGTs are malignant [6]. Similar to previous studies, the rate of dogs with malignant tumors was 40.0% (24/60) in the present study. One study reported that the risk for incidence of malignant tumors increases with increasing tumor size, in the study, the mean size of benign tumors was 21 mm and that of malignant tumors was 47 mm [20]. In this study, similar results were found that the mean size of benign tumors was 20 mm, but that of malignant tumors was 48.7 mm. In addition, there was a correlation between tumor size and the incidence of malignant tumors. Only 23.7% of tumors less than 30 mm were malignant, but 50.0% of tumors between 30 mm and 50 mm, and 78.6% of tumors larger than 50 mm were malignant.

There have been no reports on the quantity of tumors related to the incidence of malignant tumors. There was no significant correlation between the number of tumors and malignancy in the present study. However, 22.2% of solitary MGTs, 33.3% of




Fig. 1. Kaplan-Meier analysis for 2-year survival rate of 60 dogs with mammary gland tumors after surgery according to (A) clinical TNM stage and (B) histopathological results. (A) The 2-year survival rate for dogs with low TNM stage was 87.8%, and 33.3% ($p < 0.001$) for dogs with high-level TNM stage. Mean survival time of dogs with low-level TNM stage was 22.2 ± 0.8 months, but for those with a high-level TNM stage was 11.2 ± 3.6 months. (B) The 2-year survival rate for dogs with benign tumors was 90.2% and for malignant tumors was 67.3% ($p = 0.008$). Mean survival of dogs with benign tumors was 23.1 ± 0.6 months, but 17.7 ± 2.0 months for dogs with malignant tumors.

double MGTs, and 45.2% of multiple MGTs were diagnosed as malignant in this study. Considering this point, the possibility of malignancy could not be ruled out even for a single MGT. This could be the basis for early wide resection rather than simple mastectomy or lumpectomy.

Several studies have revealed that increased age, large tumor size, progressed TNM stage, and malignant tumors were related to poor prognosis [8,13-18,20]. Dogs with progressed TNM stage had shorter median survival time after tumor resection compared with dogs with clinical stage I, II, or III [17]. OHE at the time of tumor resection prolonged survival time in one study [12], but not in others [9-11].

According to several studies, the most important prognostic factor for canine MGTs is tumor size [13-16]. In this study, as the tumor size increased 10 mm, the risk of death within 2 years after surgery also increased 1.213 times ($p = 0.011$). In addition, tumor size was correlated with the development of malignant MGTs. The larger the tumor size, the higher the likelihood of malignancy and metastasis, which are considered to affect the 2-year survival rate after tumor resection.

Unlike tumor size, the side and location of affected glands have been reported to be unrelated to prognosis [17]. Likewise, in this study, the number of tumors and the number of affected mammary glands did not affect the incidence of malignant tumors or the 2-year survival rate after tumor resection. Our results suggest that only tumor size was a risk factor associated with the development of malignant MGTs.

Dogs with benign MGTs and approximately 50% of dogs with malignant MGTs are treated with surgery alone [6,7]. However, dogs that previously had MGTs have a risk of developing new tumors in other mammary glands. Because approximately 58% of dogs with MGTs relapse after surgical removal of benign tumors, regional mastectomy is recommended, even with single tumors suspected to be benign [23].

In our cases, 9 dogs died due to MGTs within 2 years after tumor resection: 2 dogs had benign tumors, and one of them died because of the occurrence of inflammatory carcinoma after surgery. Another dog was diagnosed with tumor relapse and lymph node metastasis and died during chemotherapy. Except for these 9 dogs, dogs with benign tumors or without metastasis had a good prognosis.

According to the results of present study, high-level TNM stage was closely related with the incidence of malignant tumors. The larger tumor size, the more likely it is to be the malignant tumor, and the malignant tumor have the risk of metastasis. Considering these points, tumor size is one of the most important factor associated with incidence of malignancy and

2-year survival after tumor resection. Therefore, surgical intervention in the early stages is considered to be helpful for increasing the 2-year survival rate.

In conclusion, the prognostic factors were tumor size and high-level TNM stage. Among these, tumor size could be an important criterion for the preoperative determination of malignant tumors. Because tumors larger than 30 mm are highly likely to be malignant, metastatic evaluation is required, and wide resection is recommended.

Acknowledgments

This work was supported by a research fund from the Chungnam National University.

ORCID

Chang-Hwan Moon, <https://orcid.org/0000-0003-3530-9860>

Dae-Hyun Kim, <https://orcid.org/0000-0002-3525-3812>

Sung-Ho Yun, <https://orcid.org/0000-0002-9027-3859>

Hae-Beom Lee, <https://orcid.org/0000-0002-1594-6670>

Seong-Mok Jeong, <https://orcid.org/0000-0001-8653-5927>

References

1. Sleeckx N, de Rooster H, Veldhuis Kroeze EJ, Van Ginneken C, Van Brantegem L. Canine mammary tumours, an overview. *Reprod Domest Anim* 2011;46:1112-1131.
2. Dorn CR, Taylor DO, Schneider R, Hibbard HH, Klauber MR. Survey of animal neoplasms in Alameda and Contra Costa Counties, California. II. Cancer morbidity in dogs and cats from Alameda County. *J Natl Cancer Inst* 1968;40:307-318.
3. MacEwen EG, Vail DM, Page RL. Withrow & MacEwen's Small Animal Clinical Oncology. 5th ed. pp. 356-372, Elsevier Saunders, St. Louis, 2013.
4. Meuten DJ. Tumors in Domestic Animals. 4th ed. pp. 576-606, Iowa State University Press, Ames, 2002.
5. Fossum TW. Small Animal Surgery. 4th ed. pp. 809-815, Elsevier Saunders, St. Louis, 2013.
6. Misdorp W, Hart AA. Canine mammary cancer. II. Therapy and causes of death. *J Small Anim Pract* 1979;20:395-404.
7. Straw R. Treatment of Mammary Gland Tumours and Perianal Neoplasia. pp. 672-675, North American Veterinary Conference, Orlando, 2005.
8. Hellmén E, Bergström R, Holmberg L, Spångberg IB, Hansson K, Lindgren A. Prognostic factors in canine mammary

- tumors: a multivariate study of 202 consecutive cases. *Vet Pathol* 1993;30:20–27.
9. Brodey RS, Fidler IJ, Howson AE. The relationship of estrous irregularity, pseudopregnancy, and pregnancy to the development of canine mammary neoplasms. *J Am Vet Med Assoc* 1966;149:1047–1049.
 10. Misdorp W. Canine mammary tumours: protective effect of late ovariectomy and stimulating effect of progestins. *Vet Q* 1988;10:26–33.
 11. Schneider R, Dorn CR, Taylor DO. Factors influencing canine mammary cancer development and postsurgical survival. *J Natl Cancer Inst* 1969;43:1249–1261.
 12. Sorenmo KU, Shofer FS, Goldschmidt MH. Effect of spaying and timing of spaying on survival of dogs with mammary carcinoma. *J Vet Intern Med* 2000;14:266–270.
 13. Pérez Alenza MD, Peña L, Nieto AI, Castaño M. Clinical and pathological prognostic factors in canine mammary tumors. *Ann Ist Super Sanita* 1997;33:581–585.
 14. Ferreira E, Bertagnolli AC, Cavalcanti MF, Schmitt FC, Cassali GD. The relationship between tumour size and expression of prognostic markers in benign and malignant canine mammary tumours. *Vet Comp Oncol* 2009;7:230–235.
 15. Philibert JC, Snyder PW, Glickman N, Glickman LT, Knapp DW, Waters DJ. Influence of host factors on survival in dogs with malignant mammary gland tumors. *J Vet Intern Med* 2003;17:102–106.
 16. Yamagami T, Kobayashi T, Takahashi K, Sugiyama M. Prognosis for canine malignant mammary tumors based on TNM and histologic classification. *J Vet Med Sci* 1996;58:1079–1083.
 17. Chang SC, Chang CC, Chang TJ, Wong ML. Prognostic factors associated with survival two years after surgery in dogs with malignant mammary tumors: 79 cases (1998-2002). *J Am Vet Med Assoc* 2005;227:1625–1629.
 18. Sorenmo K. Canine mammary gland tumors. *Vet Clin North Am Small Anim Pract* 2003;33:573–596.
 19. Millanta F, Calandrella M, Bari G, Niccolini M, Vannozzi I, Poli A. Comparison of steroid receptor expression in normal, dysplastic, and neoplastic canine and feline mammary tissues. *Res Vet Sci* 2005;79:225–232.
 20. Sorenmo KU, Kristiansen VM, Cofone MA, Shofer FS, Breen AM, Langeland M, Mongil CM, Grondahl AM, Teige J, Goldschmidt MH. Canine mammary gland tumours; a histological continuum from benign to malignant; clinical and histopathological evidence. *Vet Comp Oncol* 2009;7:162–172.
 21. Owen LN, World Health Organization. *TNM Classification of Tumours in Domestic Animals*. World Health Organization, Geneva, 1980.
 22. MacEwen EG, Harvey HJ, Patnaik AK, Mooney S, Hayes A, Kurzman I, Hardy WD Jr. Evaluation of effects of levamisole and surgery on canine mammary cancer. *J Biol Response Mod* 1985;4:418–426.
 23. Stratmann N, Failing K, Richter A, Wehrend A. Mammary tumor recurrence in bitches after regional mastectomy. *Vet Surg* 2008;37:82–86.